

INTER-AMERICAN TROPICAL TUNA COMMISSION

WORKING GROUP TO REVIEW STOCK ASSESSMENTS

8TH MEETING

LA JOLLA, CALIFORNIA (USA)
7-11 MAY 2007

DOCUMENT SAR-8-14

**SEABIRDS: INTERACTIONS WITH LONGLINE FISHERIES: AREAS AND
MITIGATION TOOLS**

IATTC Resolution [C-05-01](#)¹ recommends that Parties implement the [FAO International Plan of Action for Reducing Incidental Catches of Seabirds in Longline Fisheries](#)², report on their National Plans of Action to the Commission, and report on incidental seabird bycatches from all fisheries under the purview of the IATTC. The Stock Assessment Working Group was also asked to provide an assessment of the impact of bycatches on seabird populations and identify areas of potential interactions. Similar resolutions have been adopted by the Indian Ocean Tuna Commission (IOTC) and the Western and Central Pacific Fisheries Commission (WCPFC); both these organizations have also adopted requirements for mitigation measures. [IOTC Resolution 06/04](#)³ requires vessels operating south of 30°S to use bird-scaring lines (streamers, or tori lines, that keep seabirds away from the area where the bait is being set); other mitigation measures will be considered in 2007. [WCPFC Conservation and Management Measure 2006-02](#)⁴ requires vessels to use at least one mitigation measure from a menu of options, with at least two being required when fishing north of 23°N or south of 30°S.

This document discusses a range of potential management actions that could be taken by the IATTC to address seabird bycatch. The 6th meeting of the IATTC's Bycatch Working Group, held in February 2007, recommended that "the Stock Assessment Working Group suggest areas where mitigation measures for reducing seabird mortality could be most effectively adopted (*i.e.*, where bird distributions and longline effort overlap), as well as suggest possible mitigation measures in these areas of vulnerability. The Commission should then consider mitigation measures at its June 2007 meeting." Information on seabirds was presented in various documents prepared for the 7th meeting of the Stock Assessment Working Group in May 2006⁵. Information on vulnerable areas (areas in which the distributions of seabirds and longline effort overlapped) and mitigation gear available was summarized in [Document BWG-5-05.a.i](#)⁶, prepared for the 5th meeting of the Bycatch Working Group (Appendix 1). A recent paper by Bull (2007)⁷ provides a description of seabird mitigation measures used in longline, gillnet, and trawl fisheries, and discusses the advantages and disadvantages of each method.

1. AREAS OF VULNERABILITY

As described in Appendix 1, albatrosses and petrels are the seabird species of greatest concern, being most likely to forage from baited hooks, and having life history characteristics that make them vulnerable to bycatch mortality. Of particular concern is the waved albatross, an endemic species that nests in the Galapagos Islands and forages only in the EPO (Appendix 1, Figure 1). In addition, the IATTC Area is

¹ <http://www.iattc.org/PDFFiles2/C-05-01-Seabirds.pdf>

² http://www.fao.org/fi/website/FIRetrieveAction.do?dom=org&xml=ipoa_seabirds.xml

³ [http://www.iotc.org/files/proceedings/2006/s/IOTC-2006-S10-R\[EN\].pdf](http://www.iotc.org/files/proceedings/2006/s/IOTC-2006-S10-R[EN].pdf)

⁴ [http://www.wcpfc.int/pdf/Conservation%20and%20Management%20Measure-2006-02%20\[Seabirds\].pdf](http://www.wcpfc.int/pdf/Conservation%20and%20Management%20Measure-2006-02%20[Seabirds].pdf)

⁵ <http://www.iattc.org/PDFFiles2/SAR-7-05b-Albatross-and-petrel-distribution.pdf>

<http://www.iattc.org/PDFFiles2/SAR-7-05c-Seabirds-and-Fisheries-EPO.pdf>

<http://www.iattc.org/PDFFiles2/SAR-7-10-Seabird-interactions.pdf>

⁶ <http://www.iattc.org/PDFFiles2/BWG-5-05.a.i-Seabird-assessment.pdf>

⁷ <http://www.blackwell-synergy.com/doi/full/10.1111/j.1467-2979.2007.00234.x>

important for species which predominantly breed outside the IATTC Area, but use the area extensively for foraging. This includes Laysan and black-footed albatrosses which breed in the Northwest Hawaiian Islands (Appendix 1, Figure 2), the black-browed albatross that breeds in southern Chile, and the Chatham, Buller's, and Salvin's albatrosses, which breed in New Zealand (Appendix 1, Figures 3-5).

Areas where albatross distributions overlap with IATTC longline fishing effort are presented in Appendix 1 (industrial longliners) and Figure 1 (artisanal longliners). While industrial and artisanal fisheries appear to pose different bycatch risks for seabirds (*e.g.*, Moreno *et al.* 2006⁸), defining what is an artisanal fishery can be problematic, particularly as such fisheries become more mechanized (see Gillett 2005⁹ for a discussion). Gillett proposed two categories of artisanal longline fisheries: small-scale fisheries using undecked vessels less than 12 m in length that are unpowered or use outboard motors or sails and that make fishing trips less than three days, and medium-scale fisheries comprised of decked vessels between 12 and 24 m in length with an ice hold and that make fishing trip less than two weeks within the EEZs of the vessel's country or neighboring countries. The amount of freeboard of the vessel can be considered as well, as artisanal vessels in the EPO tend to have decks that are less than 2 m above the waterline.

2. MITIGATION MEASURES TO REDUCE BYCATCHES OF SEABIRDS

While longline fisheries can pose a significant threat to some seabird populations, the extent of the threat depends greatly on the characteristics of the vessel, the degree of mechanization of the fishing operations, the setting techniques used, and whether seabird deterrent devices are used. Gear modifications are designed to keep seabirds away from the baited hooks as they are being set and retrieved, or to make the baited hooks sink more rapidly out of reach of the birds (see review and Table 3 in [SAR-7-05c](#)¹; Appendix 1; Bull 2007⁷; Gilman *et al.* 2007¹⁰). Simple devices, such as bird-scaring lines (streamers or tori lines) or bird curtains, can keep seabirds from getting close to the bait. Side-setting deters birds from approaching the hooks until they reach the rear of the vessel and have had a chance to sink deeper underwater. Other methods, such as the use of blue-dyed bait, make it more difficult for diving seabirds to see the bait in the water. Setting and retrieving longlines only at night takes advantage of the fact that many seabirds are less active at night. Management of offal discharge can either reduce the attraction of seabirds to the boat by avoiding the discard of fish parts and garbage while the line is being set or can distract seabirds away from the longline while it is being set. Area closures have also been used to reduce bycatches.

3. MANAGEMENT AND RESEARCH OPTIONS

The following are potential options that can be adopted to reduce seabird bycatch. It should be noted that seabird bycatch reduction has been most effective when a suite of measures have been adopted, rather than just a single measure (Melvin and Baker 2006)¹¹.

3.1. Information and research needs

- 1) Basic information about how the industrial and artisanal longline fisheries currently operate is required. Data should be collected on vessel numbers, types, setting operations (*e.g.*, time of day, setting methods, gear used), target species, and gear for each industrial and artisanal longline vessel fishery under the purview of the IATTC.
- 2) Observers could collect seabird bycatch data from both industrial and artisanal longliners. The data could include species identification, disposition of bycatch, condition of bycatch upon release, collection of life-history samples and data, photographs, sightings of species of interest during set/haul, tags, and a description of the seabird deterrents used. Observer data would provide a better

⁸ <http://dx.doi.org/10.1016/j.biocon.2005.07.011>

⁹ Pages 171-189 in Bayliff *et al.* (2006): <ftp://ftp.fao.org/docrep/fao/008/y5984e/y5984e00.pdf>

¹⁰ <http://www.blackwell-synergy.com/doi/full/10.1111/j.1444-2906.2007.01321.x>

¹¹ http://www.wsg.washington.edu/pdfs/Pelagic_Workshop_Rep.pdf

estimate of the magnitude of the bycatch, a survey of the types of gear and mitigation measures used in the various fleets, and an opportunity to monitor the effectiveness of the mitigation measures. A significant observer program, run for a sufficient period of time, that demonstrated that there are no impacts caused by the operation, could preclude further mitigation measures.

- 3) Continued research for reducing seabird bycatch in pelagic longlines should be encouraged. Research directions and priorities have been proposed by seabird experts at a workshop on Seabird Bycatch Mitigation in Pelagic Longline Fisheries held in Hobart, Tasmania, in October 2006 (Melvin and Baker 2006)¹¹.

3.2. Gear and fishing operations requirements

- 1) All industrial longliners could be required to adopt measures that would reduce seabird bycatch. Following the approach of the Western and Central Pacific Fisheries Commission, all industrial longline vessels could be required to adopt at least one of the mitigation measures listed in Table 1 when fishing in the IATTC Area. At least two measures, with at least one measure from Column A, would be required when fishing north of 23°N, south of 30°S, or within the area off the coast of South America and Mexico delineated in Figure 2. It may be premature to include artisanal longline vessels in these requirements because of the lack of information about these fleets and because observer data do not show significant, if any, seabird bycatch from artisanal longline vessels

TABLE 1. Mitigation measures to reduce seabird bycatch, based on the WCPFC approach³. All industrial longline vessels could be required to adopt at least one of the mitigation measures listed below (from Column A or B) when fishing in the IATTC Area. At least two measures, with at least one measure from Column A, would be required when fishing north of 23°N, south of 30°S, or within the area off the coast of South America and Mexico delineated in Figure 1. For a fuller description of these methods see IOTC³, WCPFC⁴, Bull (2007)⁷, and Gilman *et al.* (2007)¹⁰.

Column A	
<i>Side setting with a bird curtain and weighted branch lines</i>	Side setting reduces the seabirds' ability to take the baited hooks. The addition of a bird curtain keeps the seabirds away from baited hooks, and weighted branch lines allow the gear to sink even faster.
<i>Night setting with minimum deck lighting</i>	Setting and retrieving the gear at night shifts fishing operations to a time when the seabirds are less active, and reduces the visibility of the bait.
<i>Bird-scaring lines</i>	A system of streamers is deployed to keep seabirds away from the area where the longline enters the water.
<i>Weighted branch lines</i>	Weighted lines can sink the baited hooks more rapidly.
Column B	
<i>Bird-scaring lines</i>	A system of streamers is deployed to keep seabirds away from the area where the longline enters the water.
<i>Weighted branch lines</i>	Weighted lines can sink the baited hooks more rapidly.
<i>Blue-dyed bait</i>	Coloring bait reduces contrast with the water, making it more difficult for the seabirds to see.
<i>Underwater setting devices</i>	Devices such as line shooters, underwater chutes, and deep-setting capsules may set the baited hooks deep enough to reduce the seabirds' access to the bait; several of these methods are under development.
<i>Management of offal discharge</i>	Fishermen avoid discarding offal while the longline is being set to reduce the attraction by seabirds.




FIGURE 1. Locations of longline sets by vessels from Costa Rica, Ecuador, El Salvador, Guatemala, Panama, and Peru, for which observers recorded data on the catches using different type of hooks.




FIGURE 2. Potential areas (in blue) for requiring mitigation measures for reducing seabird bycatch. The areas includes waters in the IATTC Area north of 23°N and south of 30°S, plus waters in the IATTC Area bounded by the coastline at 2°N, west to 2°N, 95°W, south to 15°S, 95°W, east to 15°S, 85°W, and south to 30°S.

Appendix 1

COMISIÓN INTERAMERICANA DEL ATÚN TROPICAL INTER-AMERICAN TROPICAL TUNA COMMISSION

WORKING GROUP ON BYCATCH

5TH MEETING

BUSAN, KOREA
24 JUNE 2006

DOCUMENT BWG-5-05.a.i

REVIEW OF SEABIRD STATUS AND INCIDENTAL CATCH IN EASTERN PACIFIC OCEAN FISHERIES

IATTC [Resolution C-05-01](#) on incidental mortality of seabirds recommends the implementation of the FAO's International Plan of Action for Reducing the Incidental Catch of Seabirds in Longline Fisheries (IPOA-Seabirds; FAO 1999); the collection of all available information on interactions with seabirds, including incidental catches in all fisheries under the purview of IATTC; and for the Working Group on Stock Assessment to assess the impact of incidental catch of seabirds resulting from the activities of all the vessels fishing for tunas and tuna-like species, in the eastern Pacific Ocean, including an identification of the geographic areas where there could be interactions between longline fisheries and seabirds. This report is based on Documents SAR-7-05c, SAR-7-05b, and SAR-7-10 presented at the meeting of the Stock Assessment Review Working Group in May 2006. In its report, the Working Group noted that it did not review these documents at the same level as it did the documents on stock assessments of tunas.

Albatrosses and petrels have been identified as some of the most vulnerable species to bycatch in fisheries (Wooller *et al.* 1992; Brothers *et al.* 1999). Albatrosses, in particular, scavenge for food, and are attracted to the baited hooks during longline setting operations. The interaction of seabirds with fisheries (particularly those using longline gear) has become an international issue as demonstrated by the adoption of the IPOA-Seabirds, resolutions by regional fishery management organizations (CCAMLR, IATTC, ICCAT, IOTC, WCPFC), and mandatory mitigation requirements by CCAMLR and CCSBT. In 2004, the [Agreement on the Conservation of Albatrosses and Petrels](#) (ACAP) entered into force; it identified fishery interactions as a key threat facing these seabirds, and recommended that collaborations with regional fishery management organizations be pursued to reduce fisheries interactions (ACAP 2004).

This report reviews the status and distribution of albatrosses and petrels that breed or forage in the IATTC Area defined under the Antigua Convention (an area extending to 50°N and 50°S, bounded to the west by 150°W), and that are of known conservation concern or known to be vulnerable to bycatch, particularly in longline fisheries. Population sizes and status of albatrosses and petrels breeding or foraging in the IATTC Area are given in Table 1.

1. SEABIRD SPECIES OF CONCERN

1.1. Waved albatross

The waved albatross breeds almost exclusively on Isla Española, Galápagos, Ecuador (Anderson *et al.* 2002). The species is listed as 'vulnerable' under the International Union for the Conservation of Nature (IUCN) criteria, based on its very small breeding range (IUCN 2004). A 2001 assessment estimated a population size of 31,000 to 35,000 adults, indicating a marked decline in breeding population size between 1994 and 2001 (Anderson *et al.* 2002).

Tracking data¹² during the April-December breeding season (Figure 1) reveal that foraging is focused in the Peruvian upwelling area between the Galapagos Islands and the coast of Peru (Anderson *et al.* 1998; 2003). IATTC observer data from the purse-seine fleet, overlaid on longline effort distribution, showed a similar distribution pattern (SAR-7-10). Few observations of waved albatross have been made outside the Peruvian upwelling area (Tickell, 2000; SAR-7-10), and sightings indicate that the highest aggregations of waved albatross on the Peruvian continental shelf occur during the non-breeding season (Goya and Cardenas, 2003), suggesting that this area is highly important for this species.

TABLE 1. Albatrosses and petrels breeding or foraging in the IATTC Area

Common	Scientific	Global annual breeding pairs ¹	Status ²
Antipodean albatross ³	<i>Diomedea antipodensis</i>	5,154	Vulnerable
Black-browed albatross	<i>Thalassarche melanophrys</i>	609,395	Endangered
Black-footed albatross	<i>Phoebastria nigripes</i>	64,617	Endangered
Buller's albatross	<i>Thalassarche bulleri</i>	31,435	Vulnerable
Chatham albatross	<i>Thalassarche eremita</i>	4,000	Critically endangered
Grey-headed albatross	<i>Thalassarche chrysostoma</i>	106,036	Vulnerable
Laysan albatross	<i>Phoebastria immutabilis</i>	554,699	Vulnerable
Light-mantled albatross	<i>Phoebetria palpebrata</i>	22,031	Near threatened
Northern royal albatross	<i>Diomedea sanfordi</i>	2,078	Endangered
Salvin's albatross	<i>Thalassarche salvini</i>	76,943	Vulnerable
Short-tailed albatross	<i>Phoebastria albatrus</i>	362	Vulnerable
Southern royal albatross	<i>Diomedea epomophora</i>	7,872	Vulnerable
Wandering albatross	<i>Diomedea exulans</i>	7,874	Vulnerable
Waved albatross	<i>Phoebastria irrorata</i>	18,210	Vulnerable
Black petrel	<i>Procellaria parkinsoni</i>	2,500	Vulnerable
Grey petrel	<i>Procellaria cineria</i>	?	Near threatened
White-chinned petrel	<i>Procellaria aequinoctialis</i>	2,350,000?	Vulnerable
Southern giant petrel	<i>Macronectes giganteus</i>	29,774	Vulnerable

¹ Number of nesting pairs is the standard unit for reporting seabird abundance, based on colony monitoring. Total population estimates are not available for all species. (Sources: SAR-7-05b, BirdLife International 2004).

² Source: IUCN 2004.

³ Including Gibson's albatross (*D. antipodensis gibsoni*).

1.2. Laysan albatross

The IUCN lists the Laysan albatross as vulnerable, based on a > 30% decline over 84 years (IUCN 2004). Virtually all (~99.9%) of the world's Laysan albatross breed in the Northwestern Hawaiian Islands, with other small breeding sites are in Japan and Mexico. The current world population is estimated at 3.4 million individuals (NMFS 2005a).

Tracked Laysan albatross from the small breeding population on Isla de Guadalupe, Mexico (350 breeding pairs, BirdLife International 2004), remained almost entirely within the IATTC Area (Figure 2). Laysan albatross tracked from Tern Island (Hawaii) and the Aleutian Islands did not forage in the IATTC Area, but data from the Tagging of Pacific Pelagics (TOPP) Program indicate that Laysan albatross do disperse into the IATTC Area during the non-breeding season (S. Shaffer, cited in SAR-7-05c).

1.3. Black-footed albatross

The IUCN lists the black-footed albatross as endangered, on the basis of a projected future decline of

¹² Tracking data presented in this report are from the Global *Procellariiform* Tracking Database and the tracking plots extracted from SAR-7-05b. Contributors to this database are acknowledged in SAR-7-05b.

more than 60% over the next 56 years (IUCN 2004). Most of the world's black-footed albatross (96%) breed in the Northwestern Hawaiian Islands, with other breeding sites in Japan and Mexico. At Midway Atoll, the December 2005 count marked the fifth consecutive year of increased numbers of black-footed albatross nests, after fairly steep declines in the 1990s (J. Klavitter, cited in SAR-07-05c). The current world population of black-footed albatross is approximately 300,000 individuals (NMFS 2005a).

Tracking data show that 36% of the black-footed albatross distribution during the breeding season was within the IATTC Area (Figure 3).

1.4. Black-browed albatross

The Chilean black-browed albatross population comprises 18% of the world's population (Birdlife International 2004). This population forages along the edge of the continental shelf, spending 19% of its time within the IATTC Area during the breeding season (October until March) and 65% during the non-breeding season, when its range extended northwards into areas of longline fishing effort (Figure 4).

1.5. Chatham albatross

The Chatham albatross is listed by IUCN as critically endangered, because it is a single, small breeding population. Tracking data for the non-breeding distribution of Chatham albatross indicate that over 50% of its at-sea time is within the IATTC Area. The Chatham albatross migrates across the Pacific at or below 40°S between January to April and then moves north along the Humboldt Current into Peruvian coastal waters, wintering in a area north of 20°S, in areas which also overlaps with longline fishing effort in the IATTC Area (Figure 5). Birds then return to their breeding colony between July and September, following a more northerly route (Robertson *et al.* 2000).

1.6. Buller's albatross and Salvin's albatross

Buller's and Salvin's albatrosses are both listed as vulnerable under IUCN criteria. Both species breed in New Zealand, yet commonly forage off the South American coast (Jehl 1973; Stahl *et al.* 1998; Robertson *et al.* 2003; Spear *et al.* 2003; Goya and Cardenas 2003). Most albatrosses observed from ship surveys were within 200 km of shore (Spear *et al.* 2003). Salvin's albatross are abundant between 10 and 40°S, and Buller's albatross are most abundant below 30-40°S (Jehl 1973; Stahl *et al.* 1998, Spear *et al.* 2003). These observations indicate some likely spatial overlap with longline fisheries.

2. OTHER SEABIRDS OF POTENTIAL CONCERN

2.1. Short-tailed albatross

The short-tailed albatross breeds exclusively in Japan. As of 2005, 80-85% of the known breeding population uses a single colony on Torishima, an active volcanic island. The IUCN lists the species as vulnerable because it has a very small population (approximately 2,000 individuals) and a limited breeding range (IUCN 2004; USFWS 2005; P. Sievert, cited in SAR-7-05c). Conservation efforts have resulted in a steady population increase (7.3% per annum) and an improvement in its conservation status (IUCN 2004). It is listed here because of the species' vulnerable status and because it is uncertain how extensively its range overlaps with the IATTC Area.

2.2. Black (or Parkinson's) petrel

The black petrel breeds exclusively on two islands off New Zealand (BirdLife International 2006). The species is listed as vulnerable under the IUCN criteria, because of its limited breeding range and the potential threat of introduced predators at the colonies (IUCN 2004). The world population numbers about 10,000 individuals and is believed to be stable (IUCN 2004). It is listed here because black petrels migrate from breeding sites off New Zealand to the eastern Pacific Ocean between the Galapagos Islands, southern Mexico, and northern Peru (Pitman and Ballance 1992) and have been documented as bycatch by New Zealand longliners (Brothers *et al.* 1999).

2.3. Antipodean, southern royal, and northern royal albatrosses

These species also migrate across the Pacific from their breeding grounds in New Zealand to the coast of South America, where they remain largely south of 40°S, and generally forage south of longline fishing areas.

3. SEABIRD BYCATCH IN TUNA AND SWORDFISH FISHERIES

3.1. EPO tuna purse-seine fishery

IATTC observers are not required to record seabird mortalities during purse-seine sets because staff and observer experience indicates that they occur exceedingly rarely, if at all.

3.2. Industrial longline fisheries

Seabirds can comprise significant bycatches in longline fisheries (Melvin and Parrish 2001; Brothers *et al.* 1999). However, there may be a large variability in bycatch depending on the vessel size and setting technique and fishing area. Techniques that result in rapid line sinkage close to the hull, for example, limit the exposure of the baited hooks to seabirds and reduce bycatch.

Information on longline bycatch in the North-central and Northeast Pacific comes from the US pelagic longline fishery (SAR-7-05c). The observed fishing effort in the Hawaii deep-set tuna fishery ranged from 40°N to 0° and 180° to 135°W, and the shallow-set swordfish fishery ranged from 40°N to 15°N and 180° to 135°W, overlapping with the IATTC Area. Historically, higher levels of seabird bycatch occur in the swordfish fishery than the tuna fishery. The predominant seabird species taken in these fisheries are the black-footed and Laysan albatrosses. The estimated level of albatross bycatch has decreased since 1999 and 2000, reflecting increased use of bycatch mitigation measures, and seabird mitigation measures have been required in these fisheries since 2002.

The Hawaii-based pelagic tuna longline fleet was estimated to have taken approximately 0.004 albatross per 1,000 hooks in 2005, while the shallow-set swordfish fishery, which has 100% observer coverage, took approximately 0.04 albatross per 1,000 hooks. The total estimated 2005 take for both Hawaii fisheries was 89 black-footed and 105 Laysan albatrosses.

During 2001-2004, NMFS observers monitored the US West Coast pelagic longline swordfish fleet in an area that overlapped with the IATTC Area (SAR-7-05c), and reported 65 black-footed and 7 Laysan albatross mortalities, and 7 black-footed albatross released injured (a take rate of 0.23 birds per 1,000 hooks) (Petersen *et al.* 2003; L. Enriquez, cited in SAR-7-05c). Seabird mitigation measures were not required during this time period.

Using known bycatch rates from the US pelagic fishery based in Hawaii (1994-2000 data) and the spatial distribution of fishing effort by Japanese and Chinese Taipei fleets operating in the central North Pacific, it was estimated that 5,000 to 14,000 black-footed albatross could be taken annually in the Northeast Pacific (Lewison and Crowder 2003). However, the assumption that the Hawaii fleet and the Japanese and Chinese Taipei fleets have similar bycatch rates may not be valid (NMFS 2005c), a view shared by the IATTC staff.

Interviews with experts in the regional fisheries and seabirds suggested that there are generally low levels of bycatch of seabirds in pelagic longline fisheries in the tropical Pacific (Watling 2002).

SAR-7-05e, the report of an observer on a Chinese longline vessel that deployed 300,000 hooks, reported mortality of fewer than six birds (storm petrels and blue-footed boobies).

Integrated population modeling is being developed for black-footed and Laysan albatross populations to assess whether past and present levels of bycatch are likely to significantly affect populations. One project will attempt to estimate a threshold level of bycatch that would, at a specified level of certainty, allow a sustainable growth rate for the albatross populations (Goodman and Lebreton, 2005). Another project will

apply integrated modeling techniques to black-footed albatross populations (Maunder and Hoyle 2005).

3.3. Artisanal longline fisheries

Reports to IATTC staff from observers placed aboard artisanal longliners in the EPO by regional sea-turtle programs have not confirmed any albatross mortalities and indicate that seabird mortalities were minimal (less than a dozen seabird mortalities per ~1 million hooks set). Setting techniques will likely determine the extent of the seabird bycatch. Small coastal boats that use side-setting techniques which result in rapid line sinkage close to the hull are likely to have lower bycatches because the seabirds have limited exposure to the bait.

However, evidence exists of bycatch of albatrosses off South America (Goya and Cardenas 2003; G. Merlen, cited in Anderson *et al.* 2003; J. Awkerman *et al.*, cited in SAR-7-05b,c). A study based on 29 fisherman interviews estimated that the Peruvian artisanal longline fleet may take 2,370 to 5,610 albatrosses annually, most likely waved or Chatham albatrosses (Jahncke *et al.* 2001). In conjunction with a mark-recapture study of waved albatross, a survey of 37 major fishing communities in Perú was initiated and a minimum fishery mortality rate of 0.0093 was estimated from the tag returns (J. Awkerman *et al.*, cited in SAR-7-05b,c).

4. INDIRECT FISHERY EFFECTS ON SEABIRDS

Fisheries that target particular species can impact other components of the ecosystem, particularly those species that depend on associations with the harvested target species. Distinct seabird species assemblages form feeding flocks in association with tunas and dolphins (Au and Pitman 1986). Tropical seabirds depend upon these predators to drive prey closer to the surface (Ballance *et al.* 1997); this constitutes the most important foraging strategy for many tropical seabirds. This degree of dependence has not been found in non-tropical seabirds (Ballance and Pitman 1999). Activities that directly impact tunas can also indirectly affect the seabirds that closely associate with, and depend upon, the tunas (SAR-7-05c).

5. SEABIRD BYCATCH REDUCTION

5.1. Gear and operational changes

In the past 10 years, effective bycatch mitigation measures have been developed and used in longline fisheries (reviewed in SAR-7-05c). Gear research has addressed an array of measures and methods; recent tests (Boggs 2001; Minami and Kiyota 2002; Gilman *et al.* 2003; 2006) of gear to reduce seabird bycatch (underwater setting chute, blue-dyed bait, side-setting) have resulted in regulatory changes for the Hawaii-based pelagic longline fleet and a technical assistance program to convert vessels to side-setting operations (Brothers and Gilman 2006). When setting north of 23°N (tuna and swordfish) or shallow-setting south of 23°N, vessels must either side-set or, if the lines are set over the stern, use other methods to keep seabirds away from the baited hooks and thus avoid catching them. Vessels engaged in deep-setting using a monofilament main longline north of 23°N that do not side-set must also employ a line shooter and attach a weight of at least 45 g to each branch line within 1 m of the hook. In addition to the above requirements, shallow-setting vessels that do not side-set must also deploy gear at least 1 hour after local sunset and complete the deployment no later than local sunrise.

In the US, 44 out of 124 Hawaii longline vessels are currently side-setting, and no albatrosses were caught in 2005 by these vessels. It is not known how many vessels in non-US fleets have converted to side-setting as a means to reduce the bycatch of seabirds. Japan indicated plans for side-setting experiments on larger pelagic vessels (ISC 2006).

5.2. Area closures

The US West Coast pelagic longline fishery is prohibited in the US EEZ. Shallow sets are essentially prohibited west of 150°W. Deep sets for tuna may occur north of 23°N, provided that required seabird avoidance measures are used. Although instituted as sea turtle protective measures, these closures have

also served to reduce seabird bycatch.

5.3. Pelagic gear mitigation studies

Several promising gear devices and operational methods have been identified by fishermen and mitigation scientists that may be very effective at reducing seabird bycatch on pelagic longline gear. Studies are underway to better define an effective and practicable suite of seabird mitigation measures that can be widely applied on pelagic longline vessels. Results from these studies are anticipated within the next one to three years. International studies to test gear designed to sink baited hooks quickly such that it is inaccessible to seabirds include methods to set the line underwater or that scare the birds away from the baited hooks (SAR-7-05c). Further tests have been identified to explore methods that would make side-setting more practicable (Brothers and Gilman, 2006).

In summary:

1. Information indicates that longline fisheries in the IATTC Area may have both direct and indirect impacts on some seabird populations. The level of the impact is currently not known.
2. Remote-tracking data and at-sea observations highlight the importance of the IATTC Area for foraging and breeding of waved and Laysan albatrosses, foraging of black-footed and black-browed albatrosses, and several other albatross species from New Zealand which migrate across the Pacific to forage in the Humboldt Current.
3. Observer data from US pelagic longline fisheries indicate bycatch of Laysan and black-footed albatrosses in the Northeast Pacific. No comparable data exist from industrial longline fleets in the central and southeast Pacific.
4. Plots of seabird distributions overlaid on pelagic longline effort revealed several areas of potential vulnerability to bycatch: 1) the area between the Galápagos Islands and mainland Ecuador and Perú where waved albatross breed and forage; 2) the area north of 20°N latitude where Laysan and black-footed albatross breed and forage, particularly near the small breeding colonies off Baja California; 3) the coastal zone along South America where several species are known to forage.
5. Seabird bycatch mitigation measures have been developed which have effectively reduced seabird bycatch in longline fisheries, and more gear research is ongoing.

REFERENCES

- ACAP. 2004. Text of the Agreement on the Conservation of Albatrosses and Petrels.
http://www.acap.aq/index.php/acap/text_of_the_agreement
- Anderson, D., A. Schwandt, and H. Douglass, 1998. Foraging ranges of waved albatrosses in the eastern Tropical Pacific Ocean. Pages 180-185 *in* G. Robertson and R. Gales (eds) Albatross Biology and Conservation. Surrey Beatty and Sons, Chipping Norton, Australia.
- Anderson, D.J., K.P. Huyvaert, V. Apanius, H. Townsend, C.L. Gillikin, L.D Hill, F. Juola, E.T. Porter, D.R. Wood, C. Loughheed, and H. Vargas. 2002. Population size and trends of the waved albatross *Phoebastria irrorata*. Marine Ornithology 30:63-69.
- Anderson D.J., K.P. Huyvaert, D.R. Wood, C.L. Gillikin, B.J. Frost, and H. Mouritsen. 2003. At-sea distribution of waved albatrosses and the Galapagos Marine Reserve. Biological Conservation 110:367-373.
- Au, D.W.K., and R.L. Pitman. 1986. Seabird interactions with tuna and dolphins in the eastern tropical Pacific. Condor 88:304-317.
- Ballance, L.T., and R.L. Pitman. 1999. Foraging ecology of tropical seabirds. Pages 2057-2071 *in* N. Adams and R. Slotow (eds) Proceedings of the 22nd International Ornithological Congress, Durban. BirdLife South Africa, Johannesburg.

- Ballance, L.T., R.L. Pitman, and S.B. Reilly. 1997. Seabird community structure along a productivity gradient: importance of competition and energetic constraint. *Ecology* 78:1502-1518.
- BirdLife International 2004. Tracking Ocean Wanderers: The Global Distribution of Albatrosses and Petrels. Results from the Global Procellariiform Tracking Workshop, 1-5 September 2003, Gordon's Bay, South Africa. Cambridge, UK, BirdLife International. 100 pp.
<http://www.birdlife.org/action/science/species/seabirds/tracking.html>
- BirdLife International. 2006. Species factsheet: *Procellaria parkinsoni*. <http://www.birdlife.org>
- Boggs, C.H. 2001. Deterring albatrosses from contacting baits during swordfish longline sets. Pages 79-94. *in* E. Melvin and J.K. Parrish (eds) *Seabird Bycatch: Trends, Roadblocks, and Solutions*. University of Alaska Sea Grant, Fairbanks, Alaska.
- Brothers, N.P., J. Cooper, and S. Lokkeborg. 1999. The incidental catch of seabirds by longline fisheries: Worldwide review and technical guidelines for mitigation. FAO Fisheries Circular No. 937, FIIT/C937. FAO, Rome. 101 pp.
http://www.fao.org/documents/show_cdr.asp?url_file=/docrep/005/w9817e/w9817e00.htm
- Brothers, N., and E. Gilman. 2006. Technical assistance for Hawaii pelagic longline vessels to change deck design and fishing practices to side set. February 2006. Report for the Hawaii Longline Association, US NOAA Fisheries Pacific Islands fisheries Center and Pacific Islands Regional Office, and Western Pacific Regional Fishery Management Council. 86 pp.
<http://www.fakr.noaa.gov/protectedresources/seabirds/sidesetassistancefinal.pdf>
- FAO. 1999. The International Plan of Action for Reducing the Incidental Catch of Seabirds in Longline Fisheries. 9 pp.
http://www.fao.org/figis/servlet/static?xml=ipoa_seabirds.xml&dom=org&xp_nav=1
- Gilman, E., C. Boggs, and N. Brothers. 2003. Performance assessment of an underwater setting chute to mitigate seabird bycatch in the Hawaii pelagic longline tuna fishery. *Ocean and Coastal Management* 46:985-1010.
- Gilman, E., N. Brothers, and D.R. Kobayashi. 2006. Comparison of three seabird bycatch avoidance methods in Hawaii pelagic longline fisheries. *Fisheries Science* In press.
- Goodman, D., and J.-D. Lebreton. 2005. Integrated modeling for Hawaiian albatross populations. Progress reports submitted to Joint Institute for Marine and Atmospheric Research (JIMAR), School of Ocean and Earth Science and Technology, University of Hawaii at Manoa, Honolulu, Hawaii. 4 pp. http://www.soest.hawaii.edu/PFRP/protected_species/goodman.html
- Goya, E., and G. Cardenas. 2003. Longline fisheries and seabirds in Peru. Pages 14-15 *in* S. Lokkeborg and W. Thiele (eds) *Report of the FAO/Birdlife South American Workshop on Implementation of NPOA-Seabirds and Conservation of Albatrosses and Petrels*, Valdivia, Chile, 2-6 December 2003. FAO Fisheries Report No. 751 FIIT/R751 (En). FAO, Rome.
<ftp://ftp.fao.org/docrep/fao/007/y5742e/y5742e00.pdf>
- ISC. 2006. Report of the Sixth Meeting of the International Scientific Committee for Tuna and Tuna-Like Species in the North Pacific Ocean. Plenary Session, March 23-27, 2006. Annex 9: Report of the Bycatch Working Group Meeting, March 20-22, 2006, La Jolla, California, USA.
- IUCN. 2004. International Union for the Conservation of Nature (IUCN) Red List of Threatened Species, 2004. http://www.redlist.org/info/categories_criteria
- Jahncke, J., E. Goya, and A. Guillen. 2001. Seabird by-catch in small-scale longline fisheries in Northern Peru. *Waterbirds* 24:137-141.
- Jehl, J.R. 1973. The distribution of marine birds in Chilean waters in winter. *Auk* 90: 114-135.
- Lewis, R.L., and L.B. Crowder. 2003. Estimating fishery bycatch and effects on a vulnerable seabird population. *Ecol. Appl.* 13:743-753.

- Maunder, M., and S. Hoyle. 2005. A generally Bayesian integrated population dynamics model for protected species. Progress report submitted to Joint Institute for Marine and Atmospheric Research (JIMAR), School of Ocean and Earth Science and Technology, University of Hawaii at Manoa, Honolulu, Hawaii. 5 pp. http://www.soest.hawaii.edu/PFRP/protected_species/maunder.html
- Melvin, E.F., and J.K. Parrish (eds.). 2001. Seabird Bycatch: Trends, Roadblocks, and Solutions. University of Alaska Sea Grant. Fairbanks, Alaska. 204 pp.
- Minami H., and M. Kiyota. 2002. Effect of blue-dyed bait on reduction of incidental take of seabirds. Report to the National Research Institute of Far Seas Fisheries, Fisheries Research Agency, Shizuoka, Japan.
- NMFS. 2005a. Final Environmental Impact Statement: Seabird Interaction Avoidance Methods under the Fishery Management Plan on Pelagics Fisheries of the Western Pacific Region and Pelagic Squid Fishery Management under the Fishery Management Plan for the Pelagics Fisheries Of The Western Pacific Region and the High Seas Fishing Compliance Act. NMFS Pacific Islands Region Office (PIRO), April 2005. <http://swr.nmfs.noaa.gov/pir/feis/feis.htm>
- NMFS. 2005b. Annual report on seabird interactions and mitigation efforts in the Hawaii-based longline fishery for calendar year 2004. Pacific Islands Regional Office. Administrative Report AR-PIR-04-05, 27 pp. http://swr.nmfs.noaa.gov/pir/2004_Seabird_Annual_Report.pdf
- NMFS 2005c. Final Environmental Impact Statement: Seabird Interaction Avoidance Methods Under the Fishery Management Plan for the Pelagic Fisheries of the Western Paacific Region and Pelagic Squid Fishery Management under the Fishery Management Plan for the Pelagic Fisheries of the Western Pacific Region and the High Seas Fishing Compliance Act. NOAA, NMFS, PIRO, Honolulu, Hawaii, April 2005.
- Petersen D., L. Enriquez, and S. Fougner. 2003. Information on incidental mortality of seabirds and other protected species in the US West Coast pelagic longline fishery. CCAMLR WG-FSA-03/39.
- Pitman, R.L., and L.T. Ballance. 1992. Parkinson's petrel distribution and foraging ecology in the eastern Pacific: Aspects of an exclusive feeding relationship with dolphins. *Condor* 94:825-835.
- Robertson, C.J.R., E Bell, and D.G. Nicholls. 2000. The Chatham Albatross (*Thalassarche eremita*): At home and abroad. *Notornis* 47: 174.
- Robertson, C.J.R., E.A. Bell, N. Sinclair, and B.D. Bell. 2003. Distribution of seabirds from New Zealand that overlap with fisheries worldwide. *Science for Conservation* 233. Department of Conservation, New Zealand. 102 pp. <http://www.doc.govt.nz/Publications/004~Science-and-Research/Science-for-Conservation/PDF/sfc233.pdf>
- Spear, L.R., D.G. Ainley, and S.W. Webb. 2003. Distribution, abundance and behaviour of Buller's, Chatham Island and Salvin's Albatrosses off Chile and Peru. *Ibis* 145: 253-269.
- Stahl, J.C., J.A. Bartle, N.G. Cheshire, C. Petyt, and P.M. Sagar. 1998. Distribution and movements of Buller's albatross (*Diomedea bulleri*) in Australian seas. *New Zealand Journal of Zoology* 25:109-137.
- Tickell, W.L.N. 2000. Albatrosses. Pica Press, Sussex, UK. 448 pp.
- USFWS. 2005. Short-tailed Albatross Draft Recovery Plan. Anchorage, AK. 62 pp. http://ecos.fws.gov/docs/recovery_plans/2005/051027.pdf
- Watling, D. 2002. Interactions between seabirds and Pacific Islands' Fisheries, particularly the tuna fisheries. Report for the Secretariat of the Pacific Community by Environmental Consultants, Fiji.. 37 pp. <http://www.spc.int/coastfish/Reports/HOF3/E-IP10-HOF3.pdf>
- Wooller, R.D., J.S. Bradley, and J.P. Croxall, J.P. 1992. Long-term population studies of seabirds. *Trends in Ecology and Evolution* 7:111-114.




FIGURE 1. Distributions of waved albatrosses during the breeding season, tracked from Isla Española (>99% population), and longline fishing effort in the IATTC Area, 1997-2004 (hooks set per 5° square). Data from D. Anderson and J. Awkerman, Wake Forest University; Source: SAR-7-05b.

FIGURA 1. Distribuciones de albatros de las Galápagos durante la temporada de cría, rastreados desde Isla Española (>99% de la población), y esfuerzo de pesca palangrero en el Área de la CIAT, 1997-2004 (anzuelos calados por cuadrícula de 5°). Datos de D. Anderson y J. Awkerman, Wake Forest University; Fuente: SAR-7-05b.




FIGURE 2. Distribution of Laysan albatrosses, tracked during the breeding season from Tern Island in Hawaii, and Isla de Guadalupe in Mexico, and overlap with the IATTC Area and longline fishing effort in the IATTC Area, 1997-2004 (hooks set per 5° square). Data from R. Suryan, Hatfield Marine Science Center, Oregon State University; Source: SAR-7-05b.

FIGURA 2. Distribución de albatros de Laysan, rastreados durante la temporada de cría desde Tern Island (Hawaii) e Isla de Guadalupe (México), y superposición con el Área de la CIAT y el esfuerzo de pesca palangrero en el Área de la CIAT, 1997-2004 (anzuelos calados por cuadrícula de 5°). Datos de R. Suryan, Hatfield Marine Science Center, Oregon State University; Fuente: SAR-7-05b.




FIGURE 3. Distribution of black-footed albatrosses during the non-breeding season, tracked after capture at sea, and overlap with the IATTC Area and longline fishing effort in the IATTC Area, 1997-2004 (hooks set per 5° square). Data from R. Suryan, Hatfield Marine Science Center; D. Hyrenbach, Scripps Institute of Oceanography; R. Dotson, NMFS; Source: SAR-7-05b.

FIGURA 3. Distribución del albatros patinegro fuera de la temporada de cría, rastreados después de su captura en el mar, y superposición con el Área de la CIAT y el esfuerzo de pesca palangrero en el Área de la CIAT, 1997-2004 (anzuelos calados por cuadrícula de 5°). Datos de R. Suryan, Hatfield Marine Science Center; D. Hyrenbach, Scripps Institute of Oceanography; R. Dotson, NMFS; Fuente: SAR-7-05b.




FIGURE 4. Distribution of black-browed albatrosses during the non-breeding season, tracked from Islas Diego Ramirez, Chile, and overlap with the IATTC Area and longline fishing effort in the IATTC Area, 1997-2004 (hooks set per 5° square). Data from C. Robertson, Department of Conservation New Zealand, D. Nicholls, and M. Murray; Source: SAR-7-05b.

FIGURA 4. Distribución del albatros ojeroso fuera de la temporada de cría, rastreados desde las Islas Diego Ramirez (Chile), y superposición con el Área de la CIAT y el esfuerzo de pesca palangrero en el Área de la CIAT, 1997-2004 (anzuelos calados por cuadrícula de 5°). Datos de C. Robertson, Departamento de Conservación de Nueva Zelanda, D. Nicholls, y M. Murray; Fuente: SAR-7-05b.




FIGURE 5. Distribution of Chatham albatrosses during the non-breeding season, and overlap with IATTC Area and longline fishing effort in the IATTC Area, 1997-2004 (hooks set per 5° square). Data from C. Robertson, Department of Conservation New Zealand, D. Nicholls, and M. Murray; Source: SAR-7-05b.

FIGURA 5. Distribución del albatros de Chatham fuera de la temporada de cría, rastreados desde las Islas Diego Ramírez (Chile), y superposición con el Área de la CIAT y el esfuerzo de pesca palangrero en el Área de la CIAT, 1997-2004 (anzuelos calados por cuadrícula de 5°). Datos de C. Robertson, Departamento de Conservación de Nueva Zelanda, D. Nicholls, y M. Murray; Fuente: SAR-7-05b.