Review of the data for Taiwanese longline in the Pacific Ocean

Sheng-Ping Wang

National Taiwan Ocean University

Columns

Column name	Note
CTNO	Vessel number
Year	
Month	
Day	
Position (5 degree)	
NHBF	After 1994
Hooks	
Catch in number by species	
Catch in weight by species	
Temperture	Lots of mssing values
Bait	Lots of mssing values
Hook depth	Lots of mssing values
Position (1 degree)	After 1994

| Species |
|---|
| ALB |
| BET |
| YFT |
| PBF |
| SBT |
| TUN |
| SWO |
| MLS |
| BUM |
| BLM |
| BIL |
| SKJ |
| SKX |
| OTH |

Coverage rate

BET+YFT+ALB+SWO from logbooks / Total catches of BET+YFT+ALB+SWO

Efforts recorded from logbooks

Sets recorded by vessel scales

| CT	GRT
CT0	GRT < 5
CT1	5 ≦ GRT < 10
CT2	10 ≦ GRT < 20
CT3	20 ≦ GRT < 50
CT4	50 ≦ GRT < 100
CT5	100 ≦ GRT < 200
CT6	200 ≦ GRT < 500
CT7	500 ≦ GRT < 1000
CT8	1000 ≦ GRT

Effort distribution

Effort distribution

Effort distribution

Catches recorded from logbooks

Trend of NHBF

NHBF distribution


Regular set: NHBF < 10 hooks;
Deep set: 10 hooks ≤ NHBF < 15 hooks;
Ultra-deep: NHBF ≥ 15 hooks.

Catch composition distribution

Catch composition distribution

Catch composition distribution

# BET CPUE distribution

BET CPUE distribution

BET CPUE distribution

# YFT CPUE distribution

YFT CPUE distribution

YFT CPUE distribution